

76:651-657, (1997). Another study found that, compared to women who gave birth, women who had abortions had a 62% higher risk of death from all causes for at least *eight* years after their pregnancies. DC Reardon et. al., "Deaths Associated With Pregnancy Outcome: A Record Linkage Study of Low Income Women," *Southern Medical Journal* 95(8):834-41, (2002).

6. M Gissler et. al., "Pregnancy Associated Deaths in Finland 1987-1994 -- definition problems and benefits of record linkage," *Acta Obstetrica et Gynecologica Scandinavica* 76:651-657 (1997); and M. Gissler, "Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000," *European J. Public Health* 15(5):459-63 (2005).

7. Brian McQuarrie, "Guard, clinic at odds at abortion hearing," *Boston Globe*, April 16, 1999.

8. Carol Everett with Jack Shaw, *Blood Money* (Sisters, OR: Multnomah Books, 1992). See also Pamela Zekman and Pamela Warwick, "The Abortion Profiteers," *Chicago Sun Times* special reprint, Dec. 3, 1978 (originally published Nov. 12, 1978), p. 2-3, 33.

9. Julie A. Gazmararian et al., "The Relationship Between Pregnancy Intendedness and Physical Violence in Mothers of Newborns," *Obstetrics & Gynecology*, 85 :1031 (1995); Hortensia Amaro et al., "Violence During Pregnancy and Substance Use," *American Journal of Public Health*, 80: 575 (1990); and J. McFarlane et al., "Abuse During Pregnancy and Femicide: Urgent Implications for Women's Health," *Obstetrics & Gynecology*, 100: 27, 27-36 (2002).

10. "Is Your Mother's Feminism Dead? New Agenda for Women Revealed in Landmark Two-Year Study," press release from the Center for the Advancement of Women (www.advancewomen.org), June 24, 2003; and Steve Ertelt, "Pro-Abortion Poll Shows Majority of Women Are Pro-Life," *LifeNews.com* (www.lifenews.com/nat13.html), June 25, 2003.

11. See Theresa Burke, *Forbidden Grief: The Unspoken Pain of Abortion* (Springfield, IL: Acorn Books, 2000) and www.theunchoice.com.

12. "National Opinion Survey of 600 Adults Regarding Attitudes Toward a Pro-Woman/Pro-Life Agenda," proprietary poll commissioned by the Elliot Institute, conducted in Dec. 2002.

13. D. Reardon, *Abortion Malpractice* (Denton, TX: Life Dynamics, 1993)

14. JR Cougle, DC Reardon & PK Coleman, "Depression Associated With Abortion and Childbirth: A Long-Term Analysis of the NLSY Cohort," *Medical Science Monitor* 9(4):CR105-112, 2003.

15. Frank, et.al., "Induced Abortion Operations and Their Early Sequelae," *Journal of the Royal College of General Practitioners* 35(73):175-180, April 1985; Grimes and Cates, "Abortion: Methods and Complications", in *Human Reproduction*, 2nd ed., 796-813; M.A. Freedman, "Comparison of complication rates in first trimester abortions performed by physician assistants and physicians," *Am. J. Public Health* 76(5):550-554, 1986).

Elliot Institute, PO Box 7348, Springfield, IL 62791-7348. *Additional material is posted at [www.afterabortion.org](http://www.afterabortion.org)* Used with permission.

### The Evangelization Station

P.O. Box 267

Angels Camp, California 95222, USA

Telephone: 209-728-5598

E-mail: [evangelization@earthlink.net](mailto:evangelization@earthlink.net)

[www.evangelizationstation.com](http://www.evangelizationstation.com)

Pamphlet 636

## Forced Abortion in America

### Unwanted Abortions and other Risks and Human Rights Abuses

The Un-Choice:

- 64% of women reported feeling pressured to abort.1
- Most felt rushed or uncertain, yet 67% weren't counseled.1
- 79% weren't told of available resources.1
- 84% weren't sufficiently informed before abortion.1
- Pressure to abort can escalate to violence.2
- Homicide is the leading killer of pregnant women.3
- Clinics fail to screen for coercion.4
- Women nearly 4 times more likely to die after abortion.5
- Suicide rates 6 times higher after abortion.6
- 65% of women suffer trauma symptoms after abortion.1

**Unwanted Choice: Most abortions are unwanted or coerced. Many are forced.**

Most abortions are unwanted are coerced and many are forced, sometimes violently. Escalating pressure to abort can come from employers, husbands, parents, doctors, partners, profit-driven abortion businesses, landlords, friends and family, or even trusted financial, personal, school or religious guides, gatekeepers or authorities. They may be negligent in telling young or vulnerable

individuals or couples about available resources. They may misrepresent information or present false information as fact. They may threaten or blackmail.

These are not idle threats. Coercion can escalate to violence. As this report shows, women who resist abortion have been beaten, tortured and killed. Some have been forcibly injected with drugs, secretly given veterinary or other abortifacient drugs to force a miscarriage. Others were the victim of hit men hired to injure or kill the baby, the mother, or both.

Sometimes, the threats involve blackmail or loss of one's home or job, or the exploitation of those facing a crisis. In one case, an employer threatened to push his pregnant employee down the stairs if she didn't agree to have an abortion. In another case, a maternity-shop retailer fired pregnant employees. In yet another, a homeless woman was denied shelter until she agreed to an abortion. Often, such pressure comes from all sides and can escalate.

### **Coerced Choice ... Taken to the Clinic to Make Sure She Keeps the Appointment**

A former abortion clinic security guard testified before the Massachusetts legislature that women were routinely threatened and abused by the boyfriends or husbands who took them to the clinics to make sure they underwent their scheduled abortions.<sup>7</sup> Many women are also pressured by clinic staff financially rewarded for selling abortions.<sup>8</sup>

### **Forced Choice ... Threats Can Escalate to Violence or Homicide . the Leading Killer of Pregnant Women**

Many pregnant women have been killed by partners trying to prevent the birth. Simply being pregnant places women at higher risk of being attacked.<sup>9</sup> Homicide is the leading cause of death among pregnant women.<sup>3</sup> Women are aware of these risks. 92% of women surveyed list domestic violence and assault as the women's issue that is of highest concern to them.<sup>10</sup>

### **Uninformed Non-Choice ... .When I learned the truth, I can't tell you how betrayed I felt...**

- 54% were unsure of their decision, yet 67% received no counseling beforehand.<sup>1</sup>
- 84% were inadequately counseled beforehand.<sup>1</sup> 79% not told or deceived about available resources.<sup>1</sup>
- Many were misinformed by experts about fetal development, abortion alternatives or risks.<sup>11</sup>
- Many were denied essential personal, family, societal or economic support.<sup>11</sup>

### **Unsafe Choice ... Americans Concerned About Coercion and Risks; Support Research and Screening.**

Nearly half of voters believe coerced abortion is common. They believe negative effects are more common than generally reported.

They'll support candidates who advocate legislation holding abortionists liable for

failing to screen for evidence of coercion.<sup>12</sup> Nearly 80% of abortions take place in non-hospital facilities, ill-equipped for emergency care.<sup>13</sup>

- 31% had health complications afterwards.<sup>1</sup>
- 65% suffer multiple symptoms of post-traumatic stress disorder.<sup>1</sup>
- 65% higher risk of clinical depression.<sup>14</sup>
- 10% have immediate complications, some are life-threatening.<sup>15</sup>
- 3.5x higher risk of death from all causes.<sup>5</sup>
- Suicide rates are 6 times higher if women abort vs. giving birth.<sup>6</sup>

### **Elliot Institute: AfterAbortion.org Fact Sheets & Healing: TheUnChoice.com**

Learn More: *Download the special report, Forced Abortion in America, at [www.theunchoice.com/ForcedAbortions.pdf](http://www.theunchoice.com/ForcedAbortions.pdf)*

#### Citations

1. VM Rue et. al., "Induced abortion and traumatic stress: A preliminary comparison of American and Russian women," *Medical Science Monitor* 10(10): SR5-16 (2004).
2. See the [www.theunchoice.com](http://www.theunchoice.com) for further information and cases.
3. I.L. Horton and D. Cheng, "Enhanced Surveillance for Pregnancy-Associated Mortality-Maryland, 1993-1998," *JAMA* 285(11): 1455-1459 (2001); see also J. Mcfarlane et. al., "Abuse During Pregnancy and Femicide: Urgent Implications for Women's Health," *Obstetrics & Gynecology* 100: 27-36 (2002).
4. See [stopforcedabortions.com](http://stopforcedabortions.com).
5. M Gissler et. al., "Pregnancy Associated Deaths in Finland 1987-1994 -- definition problems and benefits of record linkage," *Acta Obstetrica et Gynecologica Scandinavica*